

INFORMATION PAGE			Form Approved OMB No. 0704-0188	
1a. REPORT Unclas 2a. SECURITY <b>AD-A234 411</b>			1b. RESTRICTIVE MARKINGS	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAFSAM-JA-90-16			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION USAF School of Aerospace Medicine		6b. OFFICE SYMBOL (If applicable) USAFSAM/HM	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Human Systems Division (AFSC) Brooks AFB, TX 78235-5300			7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION USAF School of Aerospace Medicine		8b. OFFICE SYMBOL (If applicable) USAFSAM/HM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Human Systems Division (AFSC) Brooks AFB, TX 78235-5300			10. SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO.	PROJECT NO. SUPT
			TASK NO. XX	WORK UNIT ACCESSION NO. HM
11. TITLE (Include Security Classification) The Use of the Pressure Cuff Test in the Diagnosis of Decompression Sickness				
12. PERSONAL AUTHOR(S) Rudge, Frederick W.; Stone, Jeffrey A.				
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day)
15. PAGE COUNT				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
06	10		Decompression sickness; Blood pressure cuff;	
06	05		Recompression; Hyperbaric	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Records at the Hyperbaric Medicine Division, USAF School of Aerospace Medicine, were reviewed to determine the utility of the pressure cuff test as an aid in the diagnosis of Type I decompression sickness (DCS). Applying local pressure with a blood pressure cuff has been described as a useful test to differentiate the pain of DCS from that of other musculoskeletal conditions. Records were reviewed from January 1985 to December 1989. During this period 179 patients were treated with recompression for extremity pain. In 87 patients, application of a blood pressure cuff to the painful area was used as a diagnostic aid. Only 53 patients with DCS, or 61%, had a positive test (relief of pain with local pressure). Results did not correlate with the rapidity of relief of symptoms during recompression. The conclusion is that the failure to respond to the application of local pressure should not be used to rule out the presence of DCS - this must be done with a test of pressure in a hyperbaric chamber.				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Frederick W. Rudge, Major, USAF, MC			22b. TELEPHONE (Include Area Code) (512) 536-3281	22c. OFFICE SYMBOL USAFSAM/HM

## TECHNICAL NOTE

# The Use of the Pressure Cuff Test in the Diagnosis of Decompression Sickness

FREDERICK W. RUDGE, M.D., and JEFFREY A. STONE, M.D.

RUDGE FW, STONE JA. *The use of the pressure cuff test in the diagnosis of decompression sickness.* Aviat. Space Environ. Med. 1991; 62:266-7.

Records at the Hyperbaric Medicine Division, United States Air Force School of Aerospace Medicine, were reviewed to determine the utility of the pressure cuff test as an aid in the diagnosis of Type I decompression sickness (DCS). Applying local pressure with a blood pressure cuff has been described as a useful test to differentiate the pain of DCS from that of other musculoskeletal conditions. Records were reviewed from January 1985 to December 1989. During this period 179 patients were treated with recompression for extremity pain. Application of a blood pressure cuff to the painful area was used as a diagnostic aid in 87 patients. Only 53 patients (61%) with DCS had a positive test (relief of pain with local pressure). Results did not correlate with the rapidity of relief of symptoms during recompression. We conclude that the failure to respond to the application of local pressure should not be used to rule out the presence of DCS—this must be done with a test of pressure in a hyperbaric chamber.

**D**ECOMPRESSION SICKNESS (DCS) is a physiologic syndrome which occurs as a result of excessive tissue nitrogen levels following exposure to changes in ambient pressure. In 1660, Robert Boyle was the first to note the harmful effects of acute changes in atmospheric pressure when he noted the formation of bubbles in the eye of a decompressed viper (1). Since then, many reports have been written on development of DCS in caisson and tunnel workers (2), divers (3), and aviators (4).

The most common manifestation of DCS is Type I DCS (the bends, joint pain only DCS). Among cases of DCS resulting from exposures in United States Air Force (USAF) altitude chambers, joint pain only has been noted in 69% (5). The pains of DCS are indistinguishable from the pains of other musculoskeletal disorders, such as inflammatory diseases or trauma. The pain of DCS is usually described as a deep seated aching

or burning pain, which may or may not change with movement. Vigorous rubbing or application of local pressure to the affected area often results in a temporary decrease in pain. The pressure cuff test is performed by placing a blood pressure cuff over the affected joint and inflating the cuff to above systolic pressure. Diminution of pain with application of the cuff, which recurs when the cuff pressure is released, is considered a positive test. To our knowledge, pain due to decompression sickness is the only pain which has been shown to decrease with application of local pressure.

This study reviews the USAF experience with the pressure cuff test as a tool to rule out DCS as the cause of limb pain following flying or diving. To our knowledge, no prior studies have been done on the sensitivity of the pressure cuff test in the diagnosis of decompression sickness.

## METHODS

A review was conducted of patient records maintained at the United States Air Force School of Aerospace Medicine, Hyperbaric Medicine Division. This division maintains files on all patients treated in USAF hyperbaric chambers, including the treatment of DCS. For this study, records were reviewed for the 5-year period from January 1985 to December 1989. During this period, 310 patients were treated for DCS, of whom 179 had extremity pain suitable for use of the pressure cuff test (i.e., pain involving the elbow, wrist, knee, or ankle). Patients with Type II DCS were included in the study if symptoms included pain in an extremity joint. Review of the narrative summaries and AF Form 361 (Chamber Reactor/Treatment Report) on these patients revealed that 87 patients had the pressure cuff test performed on the involved joint. Information obtained from each record included patient age, sex, joint affected, etiology of DCS (diving, flying, or altitude chamber), results of the pressure cuff test, treatment used, and time to resolution of symptoms with recompression therapy. Collected data were entered on a microcomputer for compilation.

From the U.S. Air Force School of Aerospace Medicine, Brooks Air Force Base, TX 78235-5301.

This manuscript was received for review in May 1990 and was accepted for publication in June 1990.

Address reprint requests to: Maj. Frederick W. Rudge, who is Chief of Medical Operations at the Jefferson C. Davis Hyperbaric Laboratory, USAFSAM/HM, Brooks AFB, TX 78235-5301.

# PRESSURE CUFF TEST & DECOMPRESSION SICKNESS—RUDGE & STONE

## RESULTS

Among the 87 patients with DCS who had the pressure cuff test, 76 (87%) were male. Altitude chamber exposure was the cause responsible for DCS in 89% of cases. The remainder of cases were due to diving (9%) or flying (2%). The mean age of individuals treated was 26.2, with a range of 18 to 56 years.

The results of the pressure cuff test, by affected joint, are shown in Table 1. The test was considered positive if the joint pain was partially or totally relieved with application of the cuff, and negative if pain was unchanged or increased. The pressure cuff test was positive in 61% of all cases. Among individual joints, the test was positive most frequently in the elbow at 71%. The variation in pressure cuff results in the different joints, however, is not statistically significant (chi-square (3 df) = 4.12,  $p = 0.249$ ). Pressure cuff results were also unrelated to sex (chi-square (df = 1) = 0.215,  $p = 0.643$ ), and to cause of DCS (chi-square (2 df) = 1.68,  $p = 0.431$ ).

The results of the pressure cuff test were also analyzed to assess its value in predicting the response to compression therapy. Complete resolution of pain occurred in every case. In patients with a positive pressure cuff test, the mean time to resolution of symptoms was 33 min. In patients with a negative test, the mean time was 29 min. The difference in these groups is not significant (one-factor ANOVA,  $p = 0.653$ ).

## DISCUSSION

Decompression sickness is a well-known hazard of diving and flying. Unfortunately, the joint pains of DCS are nonspecific, and cannot be distinguished from many other causes of joint pain. Any individual developing joint pain following exposure to significant hyperbaric or hypobaric pressures should be considered to have decompression sickness until proven otherwise by a test of pressure. To date, no other reliable means is available to diagnose or treat DCS.

Application of pressure to the affected joint has been shown to cause the joint pain of DCS to decrease. Lansing noted in 1944 that joint pain from DCS developing in altitude chamber occupants could be alleviated by inflating a sphygmomanometer cuff around the affected

joint (6). He noted that the amount of pressure required to give relief was variable. The pressure required for relief varied from 20 mm Hg to 190 mm Hg. Lansing noted that application of local pressure in 27 subjects with DCS caused total relief in 23 and partial relief in 4. In seven of the subjects, pain relief persisted after the pressure was removed.

Shilling (7) notes that the inflation of a blood pressure cuff above arterial pressure around an extremity joint, either at the surface or at depth, may relieve the pain and help confirm the diagnosis in a diver with DCS. Edmonds, in his text on diving medicine, also notes that the pressure cuff test may be of diagnostic value in patients with DCS (8). Hills uses the ability to relieve joint pain in DCS patients as evidence that the pain is caused by mechanical stretching of periarticular tissues by evolved bubbles of nitrogen (9).

In our experience with the pressure cuff test, the test is positive in only 61% of patients with DCS-related joint pain. Were the pressure cuff test used as the sole criterion to make the diagnosis of DCS, we would estimate that it would miss 39% of cases, which makes this test unacceptable for this purpose. Because records are not available on patients having the pressure cuff test who did not have DCS-related joint pain, no statement can be made on the specificity of the test. However, because the pressure cuff test would not be expected to be positive in a patient without DCS, the specificity of the test would be expected to be high. Therefore, a positive test should be considered as useful to confirm a diagnosis of DCS, but a negative test should not be used as a means to rule out DCS. The results of the pressure cuff test do not correlate with the speed of resolution of symptoms during recompression, and are, therefore, of no prognostic value in this regard.

## REFERENCES

- Boyle R. New experiments physico-mechanical touching the spring of the air and its effects (made for the most part, in a new pneumatic engine). Oxford: H. Hall, 1660.
- Golding FC, Griffiths PD, Hempleman HV, Paton WD, Walder DN. Decompression sickness during the construction of the Dartford Tunnel. *Br. J. Ind. Med.* 1960; 17:167-80.
- Hoiberg A. Consequences of U.S. Navy diving mishaps: decompression sickness. *Undersea Biomed. Res.* 1986; 13:383-94.
- Davis JC, Sheffield PJ, Schuknecht L, Heimbach RD, Dunn JM, Douglas G, Anderson GK. Altitude decompression sickness: hyperbaric therapy results in 145 cases. *Aviat. Space Environ. Med.* 1977; 48:722-30.
- Neubauer JC, Dixon JP, Herndon CM. Fatal pulmonary decompression sickness: a case report. *Aviat. Space Environ. Med.* 1988; 59:1181-4.
- Lansing AI. Treatment of aeroembolism by pressure application to arteries. *Air Surg. Bull.* 1944; 1:5.
- Goad RF. Diagnosis and treatment of decompression sickness. In: Shilling CW, Carlston CB, Mathias RA, eds. *The physician's guide to diving medicine*. New York: Plenum Press, 1984.
- Edmonds C, Lowry C, Pennefather J. *Diving and subaquatic medicine*. Sydney, Australia: Diving Medical Center, 1983:152.
- Hills BA. *Decompression sickness. Volume 1. The biophysical basis of prevention and treatment*. New York: John Wiley and Sons, 1977:58.

TABLE 1. RESULTS OF THE PRESSURE CUFF TEST.

Joint involved	Number of cases	Number positive (%)	Number negative (%)
Knee	55	34 (61%)	21 (39%)
Elbow	21	15 (71%)	6 (29%)
Ankle	7	3 (43%)	4 (57%)
Wrist	4	1 (25%)	3 (75%)
Total	87	53 (61%)	34 (39%)

Accession For	
HLIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	29

